

METHODS OF REDUCING PAPILLOMAVIRUS INFECTION USING IMMUNOMODULATORY POLYNUCLEOTIDE SEQUENCES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the priority benefit of U.S. Provisional application
60/188,265, filed March 10, 2000, which is hereby incorporated herein by reference
in its entirety.

STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

10 Experimental work described herein was performed at the National Institutes
of Health (NCI and NIAID divisions). The Government may have certain rights in
this invention.

TECHNICAL FIELD

15 This invention is in the field of immunomodulatory polynucleotides, more
particularly their use in ameliorating or preventing papillomavirus infection and/or
symptoms of papillomavirus infections.

BACKGROUND ART

20 Infection with human papillomavirus (HPV) is one of the most common
sexually transmitted diseases (STD) in the United States. Over 100 types of HPV
have been isolated and are categorized into two groups: cutaneous HPV and mucosal
HPV. Cutaneous HPVs include more than 15 types of HPVs that are associated with
different types of skin warts. Mucosal HPVs include more than 25 types of HPVs
25 whose main reservoir is the genital tract. Other sites for mucosal HPVs include the
respiratory tract and the oral cavity.

HPVs usually cause benign warts, or papillomas, that persist for several
months to years. In some cases, the growth of warts may become life-threatening, for

example, in the respiratory tract. In other cases, warts cause discomfort, pain, hoarseness of voice, perceived cosmetic flaws and may serve as a source of virus for sexual transmission of HPV.

Some types of mucosal HPVs (*e.g.*, HPV-16, HPV-18, HPV-31 and HPV-45) are strongly associated with the development of cervical cancer. Cervical cancer is the second most common cancer among women worldwide. Annually, approximately 450,000 new cases are diagnosed and almost 200,000 deaths are due to cervical cancer. Pisani et al. (1993) *Int. J. Cancer* 55: 891-903. The overall 5-year survival rate is about 60%; however, survival rates can range from 15%, 35%, 65%, to 85% depending on how advanced the cancer is when the patient is diagnosed. Murakami et al. (1999) *J. Immunother.* 22(3): 212-218.

Current clinical treatments for warts include the application of caustic agents (trichloroacetic acid, podophyllin, podofilox, etc.), cryotherapy, immune modulators (intralesional interferon- α , topical Imiquimod, etc.), surgical therapy, and/or application of an inhibitor of DNA synthesis, such as 5-fluorouracil. Other treatments for warts utilize VLP (Virus-Like Particles). VLPs are immunologically active particles comprised of L1 and L2 capsid proteins synthesized using yeast or bacterial vectors or recombinant techniques. VLP-based vaccine studies in dogs, cattle, and rabbits have had success in wart regression (Hines et. al. (1996) *Curr. Opin. Obstet. Gynecol.* 10: 15-19); however, successful results have not yet been seen with VLP-based vaccines in humans. This approach, while feasible, does not account for physiologically relevant features of an intact, infectious virion. Features such as viral surface conformation, antigenicity of viral oncogenes such as E6 and E7 not present in VLP, and other non-expressed proteins that may serve as potential antigens all provide reasonable targets for the immune system. The aforementioned methods eliminate warts temporarily but cannot guarantee that recurrences of warts will not occur. Furthermore, removing warts temporarily does not eradicate the HPV causing

the symptoms or address other aspects of infection, such as the possible progression of pre-neoplastic, HPV-infected cells to cancer.

In both humans and animals, benign papillomas caused by some types of HPV and animal papillomaviruses can progress to malignant cancers. The participation of co-factors, such as exposure to sunlight or x-irradiation, appear to be required for progression to malignancy in some HPV infections (e.g., HPV-5, HPV-6, HPV-8, and/or HPV-11). In animals, co-factors can include exposure to bracken fern (cattle), sunlight (sheep) and coal tar (rabbits). Co-factors that may may be associated with transformation of pre-neoplastic, HPV-infected cells to cervical cancer are still unknown but may include smoking, oral contraceptive use, pregnancy, other STDs, nutrition, immune status, and major histocompatibility complex (MHC) haplotypes.

Over 90% of cervical cancer are associated with high-grade HPV, such as HPV-16, HPV-18, and HPV-31. Bosch et al. (1995) *J. Natl. Cancer Inst.* 87:796-802. One possible cause of their high-risk potential is integration of the high-risk HPV viral genome into the host genome. The viral genome usually remains extra-chromosomal with low-risk HPV. Low-risk HPV types tend to cause more warts than high-risk HPV types and low-risk HPV types are not as strongly associated with cancer as the high-risk HPV types.

When the transformation of pre-neoplastic, HPV-infected cells to abnormal growth of cervical epithelium occurs, the abnormal growth is classified in two stages: low-grade squamous intraepithelial lesions (SIL) and high-grade SIL. Because the factors that are required for the onset of low grade SIL to occur are still unknown, a method of treatment that could prevent or delay the progression of a pre-neoplastic, HPV-infected cell to a carcinoma is highly desirable. To date, there is no known cure for cervical carcinoma. Chemotherapy is usually offered as therapy but does not guarantee that all carcinoma will be eradicated and has numerous side effects.

There is a need for a method of treatment would have one or more of the following desirable characteristics: the ability to boost anti-viral cell-mediated

immune response, preferably Th1-based response; capacity to reduce or eradicate warts in humans and animals; reduce or eradicate both HPV and animal papillomavirus infection; treat disease symptoms stemming from papillomavirus infections; prevent and/or reduce the incidence of the recurring of these symptoms; prevent and/or reduce the incidence of progression of pre-neoplastic papillomavirus-infected cells to carcinoma; be generally applicable to multiple types of human papillomavirus as well as animal papillomaviruses.

Administration of certain DNA sequences, generally known as immunostimulatory sequences or "ISS," induces an immune response with a Th1-type bias as indicated by secretion of Th1-associated cytokines. The Th1 subset of helper cells is responsible for classical cell-mediated functions such as delayed-type hypersensitivity and activation of cytotoxic T lymphocytes (CTLs), whereas the Th2 subset functions more effectively as a helper for B-cell activation. The type of immune response to an antigen is generally influenced by the cytokines produced by the cells responding to the antigen. Differences in the cytokines secreted by Th1 and Th2 cells are believed to reflect different biological functions of these two subsets. See, for example, Romagnani (2000) *Ann. Allergy Asthma Immunol.* 85:9-18.

Administration of an immunostimulatory polynucleotide with an antigen results in a Th1-type immune response to the administered antigen. Roman et al. (1997) *Nature Med.* 3:849-854. For example, mice injected intradermally with *Escherichia coli* (*E. coli*) β -galactosidase (β -Gal) in saline or in the adjuvant alum responded by producing specific IgG1 and IgE antibodies, and CD4⁺ cells that secreted IL-4 and IL-5, but not IFN- γ , demonstrating that the T cells were predominantly of the Th2 subset. However, mice injected intradermally (or with a tyne skin scratch applicator) with plasmid DNA (in saline) encoding β -Gal and containing an ISS responded by producing IgG2a antibodies and CD4⁺ cells that secreted IFN- γ , but not IL-4 and IL-5, demonstrating that the T cells were predominantly of the Th1 subset. Moreover, specific IgE production by the plasmid

DNA-injected mice was reduced 66-75%. Raz et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:5141-5145. In general, the response to naked DNA immunization is characterized by production of IL-2, TNF α and IFN- γ by antigen-stimulated CD4⁺ T cells, which is indicative of a Th1-type response. This is particularly important in treatment of allergy and asthma as shown by the decreased IgE production. The ability of immunostimulatory polynucleotides to stimulate a Th1-type immune response has been demonstrated with bacterial antigens, viral antigens and with allergens (see, for example, WO 98/55495).

Other references describing ISS include: Krieg et al. (1989) *J. Immunol.* 143:2448-2451; Tokunaga et al. (1992) *Microbiol. Immunol.* 36:55-66; Kataoka et al. (1992) *Jpn. J. Cancer Res.* 83:244-247; Yamamoto et al. (1992) *J. Immunol.* 148:4072-4076; Mojcić et al. (1993) *Clin. Immuno. and Immunopathol.* 67:130-136; Branda et al. (1993) *Biochem. Pharmacol.* 45:2037-2043; Pisetsky et al. (1994) *Life Sci.* 54(2):101-107; Yamamoto et al. (1994a) *Antisense Research and Development.* 4:119-122; Yamamoto et al. (1994b) *Jpn. J. Cancer Res.* 85:775-779; Raz et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:9519-9523; Kimura et al. (1994) *J. Biochem. (Tokyo)* 116:991-994; Krieg et al. (1995) *Nature* 374:546-549; Pisetsky et al. (1995) *Ann. N.Y. Acad. Sci.* 772:152-163; Pisetsky (1996a) *J. Immunol.* 156:421-423; Pisetsky (1996b) *Immunity* 5:303-310; Zhao et al. (1996) *Biochem. Pharmacol.* 51:173-182; Yi et al. (1996) *J. Immunol.* 156:558-564; Krieg (1996) *Trends Microbiol.* 4(2):73-76; Krieg et al. (1996) *Antisense Nucleic Acid Drug Dev.* 6:133-139; Klinman et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:2879-2883; Raz et al. (1996); Sato et al. (1996) *Science* 273:352-354; Stacey et al. (1996) *J. Immunol.* 157:2116-2122; Ballas et al. (1996) *J. Immunol.* 157:1840-1845; Branda et al. (1996) *J. Lab. Clin. Med.* 128:329-338; Sonehara et al. (1996) *J. Interferon and Cytokine Res.* 16:799-803; Klinman et al. (1997) *J. Immunol.* 158:3635-3639; Sparwasser et al. (1997) *Eur. J. Immunol.* 27:1671-1679; Roman et al. (1997); Carson et al. (1997) *J. Exp. Med.* 186:1621-1622; Chace et al. (1997) *Clin. Immunol. and Immunopathol.*

84:185-193; Chu et al. (1997) *J. Exp. Med.* 186:1623-1631; Lipford et al. (1997a) *Eur. J. Immunol.* 27:2340-2344; Lipford et al. (1997b) *Eur. J. Immunol.* 27:3420-3426; Weiner et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:10833-10837; Macfarlane et al. (1997) *Immunology* 91:586-593; Schwartz et al. (1997) *J. Clin. Invest.* 100:68-73; Stein et al. (1997) *Antisense Technology*, Ch. 11 pp. 241-264, C. Lichtenstein and W. Nellen, Eds., IRL Press; Wooldridge et al. (1997) *Blood* 89:2994-2998; Leclerc et al. (1997) *Cell. Immunol.* 179:97-106; Kline et al. (1997) *J. Invest. Med.* 45(3):282A; Yi et al. (1998a) *J. Immunol.* 160:1240-1245; Yi et al. (1998b) *J. Immunol.* 160:4755-4761; Yi et al. (1998c) *J. Immunol.* 160:5898-5906; Yi et al. (1998d) *J. Immunol.* 161:4493-4497; Krieg (1998) *Applied Antisense Oligonucleotide Technology* Ch. 24, pp. 431-448, C.A. Stein and A.M. Krieg, Eds., Wiley-Liss, Inc.; Krieg et al. (1998a) *Trends Microbiol.* 6:23-27; Krieg et al. (1998b) *J. Immunol.* 161:2428-2434; Krieg et al. (1998c) *Proc. Natl. Acad. Sci. USA* 95:12631-12636; Spiegelberg et al. (1998) *Allergy* 53(45S):93-97; Horner et al. (1998) *Cell Immunol.* 190:77-82; Jakob et al. (1998) *J. Immunol.* 161:3042-3049; Redford et al. (1998) *J. Immunol.* 161:3930-3935; Weeratna et al. (1998) *Antisense & Nucleic Acid Drug Development* 8:351-356; McCluskie et al. (1998) *J. Immunol.* 161(9):4463-4466; Gramzinski et al. (1998) *Mol. Med.* 4:109-118; Liu et al. (1998) *Blood* 92:3730-3736; Moldoveanu et al. (1998) *Vaccine* 16: 1216-1224; Brazolot Milan et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:15553-15558; Broide et al. (1998) *J. Immunol.* 161:7054-7062; Broide et al. (1999) *Int. Arch. Allergy Immunol.* 118:453-456; Kovarik et al. (1999) *J. Immunol.* 162:1611-1617; Spiegelberg et al. (1999) *Pediatr. Pulmonol. Suppl.* 18:118-121; Martin-Orozco et al. (1999) *Int. Immunol.* 11:1111-1118; EP 468,520; WO 96/02555; WO 97/28259; WO 98/16247; WO 98/18810; WO 98/37919; WO 98/40100; WO 98/52581; WO 98/55495; WO 98/55609 and WO 99/11275. See also Elkins et al. (1999) *J. Immunol.* 162:2291-2298, WO 98/52962, WO 99/33488, WO 99/33868, WO 99/51259 and WO 99/62923. See also Zimmermann et al. (1998) *J. Immunol.* 160:3627-3630; Krieg (1999) *Trends Microbiol.* 7:64-65; U.S. Patent Nos.

5,663,153, 5,723,335, 5,849,719 and 6,174,872. See also WO 99/56755, WO 00/06588, WO 00/16804; WO 00/21556; WO 00/67023 and WO 01/12223.

There remains a serious need to develop effective therapies and preventive strategies for papillomaviruses.

5 All publications and patent applications cited herein are hereby incorporated by reference in their entirety.

DISCLOSURE OF THE INVENTION

10 The invention provides methods of suppressing, ameliorating and/or preventing papillomavirus infection in an individual using immunostimulatory polynucleotide sequences. Accordingly, in one aspect, the invention provides methods of palliating, ameliorating, reducing severity, and/or eliminating one or more symptoms of *papillomavirinae* infection without administering *papillomavirinae* antigen. A polynucleotide comprising an immunostimulatory sequence (an "ISS") is
15 administered to an individual who is at risk of being exposed to *papillomavirinae*, has been exposed to *papillomavirinae* and/or is infected with *papillomavirinae*. The ISS-containing polynucleotide is administered without any *papillomavirinae* antigens. Administration of the ISS results in reduced incidence, recurrence, and/or severity of one or more symptoms of *papillomavirinae* infection.

20 In one embodiment, the invention provides methods for preventing a symptom of *papillomavirinae* infection in an individual at risk of being exposed to *papillomavirinae* which entail administering an effective amount of a composition comprising a polynucleotide comprising an immunostimulatory sequence (ISS) (*i.e.*, an amount of the composition sufficient to prevent a symptom of *papillomavirinae*
25 infection) to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition (*i.e.*, antigen is

not administered with the ISS-containing polynucleotide), thereby preventing a symptom of *papillomavirinae* infection.

In another embodiment, the invention provides methods for preventing a symptom of *papillomavirinae* infection in an individual which entail administering an effective amount of a composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby preventing a symptom of *papillomavirinae* infection. The individual may have been exposed to or infected by *papillomavirinae*.

In another embodiment, the invention provides methods for suppressing a *papillomavirinae* infection in an individual which entail administering an effective amount of a composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby suppressing a *papillomavirinae* infection. The individual may have been exposed to or infected by *papillomavirinae*.

Another embodiment of the invention provides methods of reducing severity of a symptom of *papillomavirinae* infection in an individual which entail administering an effective amount of a composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby reducing severity of a symptom of *papillomavirinae* infection. The individual may have been exposed to or infected by *papillomavirinae*.

Another embodiment of the invention provides methods of delaying development of a *papillomavirinae* infection and/or a symptom of *papillomavirinae* infection in an individual which entail administering an effective amount of a

composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby delaying development of a
5 *papillomavirinae* infection and/or a symptom of *papillomavirinae* infection. The individual may have been exposed to or infected by *papillomavirinae*.

Another embodiment of the invention provides methods of reducing duration of a *papillomavirinae* infection in an individual which entail administering an effective amount of a composition comprising a polynucleotide comprising an ISS to
10 the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby reducing duration of a *papillomavirinae* infection. The individual may have been exposed to or infected by *papillomavirinae*.

15 Another embodiment of the invention provides methods of reducing recurrence of a symptom of *papillomavirinae* infection in an individual infected with *papillomavirinae* which entail administering an effective amount of a composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a
20 *papillomavirinae* antigen is not administered in conjunction with administration of the composition, thereby reducing recurrence of a symptom of *papillomavirinae* infection.

In another aspect, the invention provides methods for suppressing papillomavirus infection in a papillomavirus-infected individual which entail
25 administering an effective amount of a composition comprising a polynucleotide comprising an ISS to the individual, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein a *papillomavirinae* antigen is not

administered in conjunction with administration of the composition, thereby suppressing papillomavirus infection.

In another aspect, the invention provides kits for use in ameliorating and/or preventing a symptom of *papillomavirinae* infection in an individual infected with, exposed to or at risk of being exposed to *papillomavirinae* and/or reduction in recurrence of a symptom of *papillomavirinae* infection. The kits comprise a composition comprising a polynucleotide comprising an ISS, wherein the ISS comprises the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' and wherein the kit does not comprise a *papillomavirinae* antigen, and wherein the kits comprise instructions for administration of the composition to an individual infected with, exposed to or at risk of being exposed to *papillomavirinae*.

In some embodiments of the methods and kits of the invention, the ISS comprises the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine, C, G-3'. In further embodiments of the methods and kits, the ISS comprises a sequence selected from the group consisting of AACGTTCG and GACGTTCG.

In some embodiments of the methods and kits of the invention, the ISS comprises the sequence 5'-TGACTGTGAACGTTCGAGATGA-3' (SEQ ID NO:1).

In some embodiments of the methods and kits of the invention, the individual is a mammal. In further embodiments, the mammal is human.

In some embodiments of the methods and kits of the invention, the *papillomavirinae* is a papillomavirus. In further embodiments of the methods and kits of the invention, the papillomavirus is a human papillomavirus (HPV) or an animal papillomavirus.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a bar graph depicting results of ISS treatment in a canine model of papillomavirus for time of wart regression.

Figure 2(A) - (D) are graphs depicting results of ISS treatment of papillomavirus in a rabbit model. The data is expressed as geometric mean diameter (GMD) over time after inoculation. Closed circles indicate Group A animals, open circles indicate Group B animals, and closed triangles indicate Group C animals. Fig. 2(A) depicts GMD for the left side, high CRPV dose lesions. Fig. 2(B) depicts GMD for the left side, low CRPV dose lesions. Fig. 2(C) depicts average GMD for the right side, high CRPV dose lesions. Fig. 2(D) depicts average GMD for the right side, low CRPV dose lesions.

Figure 3 is a graph depicting results of ISS treatment of rabbit papillomavirus. The data is expressed as geometric mean diameter (GMD) over time after inoculation. Closed circles indicate ISS treated papilloma sites, open circles indicate untreated papilloma sites animals, and downward arrows indicate timing of ISS treatments.

MODES FOR CARRYING OUT THE INVENTION

We have discovered methods of treating papillomavirus infections. The methods described herein are applicable to all *papillomavirinae* and particularly to methods of ameliorating infection (including recurrences) with a member of the *papillomavirinae* subfamily, preferably papillomavirus (including high risk and low risk). A polynucleotide comprising an immunostimulatory sequence (an "ISS") is administered to an individual who has been exposed to and/or infected with *papillomavirinae*. Administration of the ISS-containing polynucleotide without co-administration of a papillomavirus antigen results in reduced severity of one or more symptoms of *papillomavirinae* infection. It is understood that all the embodiments describes herein do not include or involve papillomavirus antigen.

The invention also relates to kits for ameliorating and/or preventing *papillomavirinae* infection and/or a symptom of *papillomavirinae* infection in exposed individuals. The kits, which do not contain an *papillomavirinae* antigen,

comprise a polynucleotide comprising an ISS and instructions describing the administration of an ISS-containing polynucleotide to an individual for the intended treatment.

5 *General techniques*

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, *Molecular Cloning: A Laboratory Manual*, second edition (Sambrook et al., 1989); *Oligonucleotide Synthesis* (M.J. Gait, ed., 1984); *Animal Cell Culture* (R.I. Freshney, ed., 1987); *Handbook of Experimental Immunology* (D.M. Weir & C.C. Blackwell, eds.); *Gene Transfer Vectors for Mammalian Cells* (J.M. Miller & M.P. Calos, eds., 1987); *Current Protocols in Molecular Biology* (F.M. Ausubel et al., eds., 1987); *PCR: The Polymerase Chain Reaction*, (Mullis et al., eds., 1994); *Current Protocols in Immunology* (J.E. Coligan et al., eds., 1991); *The Immunoassay Handbook* (David Wild, ed., Stockton Press NY, 1994); and *Methods of Immunological Analysis* (R. Masseyeff, W.H. Albert, and N.A. Staines, eds., Weinheim: VCH Verlags gesellschaft mbH, 1993).

20

Definitions

“Papillomavirus” refers to a type of virus that is in the subfamily *papillomavirinae*. Members of *papillomavirinae* are sometimes grouped within a larger family of *papovaviridae*, which includes not only the papillomaviruses, but also polyomaviruses and simian vacuolating virus. Papillomaviruses are small, nonenveloped viruses with an icosahedral symmetry, 72 capsomers, and a double-stranded DNA genome of about 8,000 base pairs. There are about ten open reading frames (ORF) and all ORFs are located on one strand. The papillomavirus genome is

25

organized into an early region and a late region. Early region genes code for proteins (E1-E8) required for viral DNA replication and cellular transformation while the late genes code for capsid protein (L1 and L2) and a regulatory region of transcriptional and replication control.

5 The term “human papillomavirus” (“HPV”) refers to papillomaviruses of human species origin and/or which are capable of infecting a human. There are over 100 types of HPV. HPV can be divided into cutaneous HPV or mucosal HPV, depending on location where the pathology and/or infection occurs. HPV can additionally be divided into “high-risk” and “low-risk”.

10 The term “high-risk” or “high-grade” refers to HPV that are strongly associated with cellular transformations that may lead to neoplasia and/or carcinoma. HPV types that are associated with development of carcinoma include, but are not limited to, HPV-16, -18, -30, -31, -33, -34, -35, -39, -40, -41, -42, -43, -44, -45, -51, -52, -56, -57, -58, -61, -62, -66, -69.

15 The term “low-risk” refers to HPV types that have lower cellular transformation potentials including, but not limited to, HPV-6 and HPV-11.

20 The term “animal papillomavirus” refers to papillomaviruses of non-human species origin, including, but is not limited to, cattle, horses, deer, rabbits, sheep, dogs, elk, nonhuman primates, rodents, harvest mice, multimammate mice, parrots and chaffinches.

 The term “papilloma” herein refers to papillomavirus-associated warts, the development, appearance, keratinous texture and histological features thereof.

 The term “condyloma” herein refers to an HPV-associated wart usually seen on the external genitalia or near the anus.

25 “Exposure” to a virus denotes encounter with virus which allows infection, such as, for example, surface to surface contact with an infected individual or tissue that results in minor trauma of basal cells and opportunity for papillomavirus to infect the traumatized basal cells, including, but are not limited to, non-barrier sexual

contact and/or intercourse with an individual with infected genitalia and abrasion of skin epithelium with infectious tissue as in the case of butcher handling infected meat products.

5 An individual is “seronegative” for a virus if antibodies specific to the virus cannot be detected in blood or serum samples from the individual using methods standard in the art, such as ELISA. Conversely, an individual is “seropositive” for a virus if antibodies specific for the virus can be detected in blood or serum samples from the individual using methods standard in the art, such as ELISA. An individual is said to “seroconvert” for a virus when antibodies to the virus can be detected in
10 blood or serum from an individual who was previously seronegative.

An individual who is “at risk of being exposed” to a papillomavirus is an individual who may encounter the virus such that the papillomavirus infects the individual (*i.e.*, virus enters cells and replicates). Because papillomaviruses are ubiquitous, generally any individual is at risk for exposure to papillomavirus. In
15 some contexts, an individual is at risk for exposure of HPV by engaging in one or more high risk behaviors, such as sexual relations without the use of barrier prophylactics with an infected individual.

A “symptom of papillomavirus infection” is any one or more symptoms of papillomavirus infection and includes, but is not limited to, the clinical presentation
20 of warts, condylomas and papillomas, all of which can be collectively referred to as “lesions”. The term “symptoms of papillomavirus infection” also includes secondary symptoms associated with warts, condylomas, papillomas and lesions. These secondary symptoms can include, but are not limited to, hoarseness of voice, breathing difficulties, pain and discomfort.

25 “Preventing a symptom of infection” by a papillomavirus means that the symptom does not appear after exposure to the virus.

“Suppressing” papillomavirus infection indicates any aspect of viral infection, such as viral replication, time course of infection, amount (titer) of virus, lesions,

and/or one or more symptoms is curtailed, inhibited, or reduced (in terms of severity and/or duration) in an individual or a population of individuals treated with an ISS-containing polynucleotide in accordance with the invention as compared to an aspect of viral infection in an individual or a population of individuals not treated in accordance with the invention. Reduction in viral titer includes, but is not limited to, elimination of the virus from an infected site or individual. Viral infection can be assessed by any means known in the art, including, but not limited to, measurement of virus particles, viral nucleic acid or viral antigens and detection of one or more symptoms of viral infection. Anti-virus antibodies are widely used to detect and monitor viral infection and generally are commercially available.

“Palliating” a disease or one or more symptoms of a disease or infection means lessening the extent and/or time course of undesirable clinical manifestations of a disease state or infection in an individual or population of individuals treated with an ISS in accordance with the invention.

As used herein, “delaying” development of a viral infection or a symptom of viral infection means to defer, hinder, slow, retard, stabilize, and/or postpone development of the disease or symptom when compared to not using the method(s) of the invention. This delay can be of varying lengths of time, depending on the history of the disease and/or individual being treated. As is evident to one skilled in the art, a sufficient or significant delay can, in effect, encompass prevention, in that the individual does not develop the disease.

“Reducing severity of a symptom” or “ameliorating a symptom” of viral infection means a lessening or improvement of one or more symptoms of viral infection as compared to not administering an ISS-containing polynucleotide.

“Reducing severity” also includes shortening or reduction in duration of a symptom. For *papillomavirinae*, these symptoms are well known in the art and include, but are not limited to, the clinical presentation of warts, condyloma and papilloma.

“Reducing duration of viral infection” means the length of time of viral infection (usually indicated by symptoms) is reduced, or shortened, as compared to not administering an ISS-containing polynucleotide.

5 “Reducing recurrence” refers to a reduction in frequency, severity and/or quantity of one or more recurrent viral symptoms in an infected individual or a population of infected individuals. When applied to a population of individuals, “reducing recurrence” means a reduction in the mean or median frequency, severity, quantity and/or duration of recurrent viral symptoms.

10 The term “infected individual” refers to an individual who has been infected with a member of *papillomavirinae*. Symptoms of *papillomavirinae* infection are well known in the art and include, but are not limited to, the clinical presentation of warts, condyloma and papilloma. “Infected individual” can include asymptomatic, infected individuals. Identification of asymptomatic, infected individuals can be accomplished by any of the biological viral detection methods known in the art, 15 which can include, but is not limited to, methods such as PCR, *in situ* hybridization and ELISA for virus-specific antibodies.

The term “eradicating papillomavirus infection” refers to elimination of the virus from the body of the infected individual. An implication from eradicating virus is that the immediate symptoms caused by the virus would also be eliminated, as well 20 as certain events or conditions associated with viral infection.

The term “low-grade squamous intraepithelial lesion” (SIL) and “high-grade squamous intraepithelial lesion” (SIL) refers to the current classification scheme to describe abnormal growth of cervical cells. The category of low-grade SIL includes HPV infection and cervical intraepithelial neoplasia (CIN) 1 while high-grade SIL 25 includes CIN 2 and 3 when the entire thickness of the epithelium is replaced by abnormal cells. This classification scheme is equivalent to a previous classification scheme in which the different stages ranged from cervical intraepithelial neoplasia (CIN) 1 (mild dysplasia) to CIN 2 (moderate dysplasia) to CIN 3 (carcinoma *in situ*).

A “biological sample” encompasses a variety of sample types obtained from an individual and can be used in a diagnostic or monitoring assay. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom, and the progeny thereof. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides. The term “biological sample” encompasses a clinical sample, and also includes cells in culture, cell supernatants, cell lysates, serum, plasma, biological fluid, and tissue samples.

“Viral titer” is a term well known in the art and indicates the amount of virus in a given biological sample. Amount of virus are indicated by various measurements, including, but not limited to, amount of viral nucleic acid; presence of viral particles; replicating units (RU); plaque forming units (PFU). Generally, for fluid samples such as blood and urine, amount of virus is determined per unit fluid, such as milliliters. For solid samples such as tissue samples, amount of virus is determined per weight unit, such as grams. Methods for determining amount of virus are known in the art and described herein.

An “individual” is a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, humans, farm animals, sport animals, rodents, primates and certain pets. Vertebrates also include, but are not limited to, birds (*i.e.*, avian individuals) and reptiles (*i.e.*, reptilian individuals).

The term “ISS” as used herein refers to polynucleotide sequences that effect a measurable immune response as measured *in vitro*, *in vivo* and/or *ex vivo*. Examples of measurable immune responses include, but are not limited to, antigen-specific antibody production, secretion of cytokines, activation or expansion of lymphocyte populations such as NK cells, CD4⁺ T lymphocytes, CD8⁺ T lymphocytes, B lymphocytes, and the like. Preferably, the ISS sequences preferentially activate a

Th1-type response. A polynucleotide for use in methods of the invention contains at least one ISS.

As used interchangeably herein, the terms "polynucleotide" and "oligonucleotide" include single-stranded DNA (ssDNA), double-stranded DNA (dsDNA), single-stranded RNA (ssRNA) and double-stranded RNA (dsRNA), modified oligonucleotides and oligonucleosides or combinations thereof. The polynucleotide can be linearly or circularly configured, or the polynucleotide can contain both linear and circular segments.

"Adjuvant" refers to a substance which, when added to an immunogenic agent such as antigen, nonspecifically enhances or potentiates an immune response to the agent in the recipient host upon exposure to the mixture.

An "effective amount" or a "sufficient amount" of a substance is an amount sufficient to effect beneficial or desired results, including clinical results. An effective amount can be administered in one or more administrations. A "therapeutically effective amount" is an amount to effect beneficial clinical results, including, but not limited to, alleviation of one or more symptoms associated with viral infection as well as prevention of disease(e.g., prevention of one or more symptoms of infection).

A microcarrier is considered "biodegradable" if it is degradable or erodable under normal mammalian physiological conditions. Generally, a microcarrier is considered biodegradable if it is degraded (*i.e.*, loses at least 5% of its mass and/or average polymer length) after a 72 hour incubation at 37° C in normal human serum. Conversely, a microcarrier is considered "nonbiodegradable" if it is not degraded or eroded under normal mammalian physiological conditions. Generally, a microcarrier is considered nonbiodegradable if it not degraded (*i.e.*, loses less than 5% of its mass and/or average polymer length) after at 72 hour incubation at 37° C in normal human serum.

The term "immunostimulatory sequence-microcarrier complex" or "ISS-MC complex" refers to a complex of an ISS-containing polynucleotide and a microcarrier. The components of the complex may be covalently or non-covalently linked. Non-covalent linkages may be mediated by any non-covalent bonding force, including by hydrophobic interaction, ionic (electrostatic) bonding, hydrogen bonds and/or van der Waals attractions. In the case of hydrophobic linkages, the linkage is generally via a hydrophobic moiety (e.g., cholesterol) covalently linked to the ISS.

As used herein, the term "comprising" and its cognates are used in their inclusive sense; that is, equivalent to the term "including" and its corresponding cognates.

As used herein, the singular form "a", "an", and "the" includes plural references unless indicated otherwise. For example, "a" symptom of viral infection includes one or more additional symptoms.

Methods of the invention

The invention provides methods for preventing one or more symptoms of papillomavirus infection, treating, reducing severity of and/or delaying development of one or more symptoms of papillomavirus infection, treating and/or eradicating papillomavirus infection, reducing recurrence of one or more symptoms of papillomavirus infection by administering an ISS-containing polynucleotide (used interchangeably herein with "ISS") without a papillomavirus antigen. Papillomavirus can be any of the members of the *papillomavirinae* subfamily, preferably one or more of the HPV types or animal papillomaviruses of any type, including high-risk or low-risk types. An ISS-containing composition which does not include papillomavirus antigen is administered to individuals who are infected with papillomavirus, who have been exposed to papillomavirus or who are at risk of being exposed to papillomavirus. Individuals receiving ISS are preferably mammal, more preferably human. In accordance with the invention, papillomavirus antigen is not administered

to the individual in conjunction with administration of an ISS (*i.e.*, is not administered in a separate administration at or about the same time of administration of the ISS).

In some embodiments, the individual is at risk of being exposed to virus.

5 Determination of an at risk individual is based on one or more factors that are associated with disease development and are generally known by, or can be assessed by, a skilled clinician. At risk individuals may be especially suitable candidates to receive ISS, as these individuals are generally considered to be particularly susceptible to developing symptoms of infection, which could also further lead to
10 other complications. For example, in the context of HPV infection, any individual is considered at risk, due to the wide spread prevalence of HPV infection. As another example, an individual at risk is an individual who practices unsafe sexual practices (*e.g.*, engages in anal-genital or genital-genital contact without the use of barrier-type prophylactics).

15 In other embodiments, the individual is, or has been exposed to and/or infected with papillomavirus. Exposure can be indicated by participation in unsafe sexual practices and/or development of one or more symptoms associated with viral infection. For example, in dogs, oral warts may be construed as one symptom stemming from infection with canine papillomavirus. In another instance, in humans,
20 skin warts may construed as one symptom stemming from infection with a type of HPV. The infected individual may or may not be symptomatic. Determination of infection can be based on any clinical indicia of infection, such as lesions or warts. Determination of infection in an asymptomatic individual can be accomplished by any of the methods known in the art, including, but not limited to, PCR techniques, *in*
25 *situ* hybridization and ELISA for papillomavirus-specific antibodies. In other embodiments, the individual, preferably a human, is infected with papillomavirus, with or without symptoms stemming from infection, and may be at a stage prior to the development of carcinoma. Infection with papillomavirus can be ascertained by

any of the aforementioned biological methods, while carcinoma may be detected with methods known in the art, including, but not limited to, Pap smears, biopsies, or other methods to observe morphological changes of epithelial cells.

5 ISS

The methods of this invention entail administering an immunomodulatory polynucleotide comprising an ISS (or a composition comprising such a polynucleotide). In accordance with the present invention, the immunomodulatory polynucleotide contains at least one ISS, and may contain multiple ISSs. The ISSs may be adjacent within the polynucleotide, or they may be separated by additional nucleotide bases within the polynucleotide. Alternately, multiple ISSs may be delivered as individual polynucleotides.

ISS have been described in the art and may be readily identified using standard assays which indicate various aspects of the immune response, such as cytokine secretion, antibody production, NK cell activation and T cell proliferation. See, *e.g.*, WO 97/28259; WO 98/16247; WO 99/11275; Krieg et al. (1995); Yamamoto et al. (1992); Ballas et al. (1996); Klinman et al. (1997); Sato et al. (1996); Pisetsky (1996a); Shimada et al. (1986) *Jpn. J. Cancer Res.* 77:808-816; Cowdery et al. (1996) *J. Immunol.* 156:4570-4575; Roman et al. (1997); and Lipford et al. (1997a).

The ISS can be of any length greater than 6 bases or base pairs and generally comprises the sequence 5'-cytosine, guanine-3', preferably greater than 15 bases or base pairs, more preferably greater than 20 bases or base pairs in length. As is well-known in the art, the cytosine of the 5'-cytosine, guanine-3' sequence is unmethylated. An ISS may also comprise the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine, C, G-3'. An ISS may also comprise the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine, C, C-3'. As indicated in polynucleotide sequences below, an ISS may comprise (*i.e.*, contain one or more of) the sequence 5'-

T, C, G-3'. In some embodiments, an ISS may comprise the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3' (such as 5'-CGTTCG-3'). In some embodiments, an ISS may comprise the sequence 5'-C, G, pyrimidine, pyrimidine, C, G, purine, purine-3'. In some embodiments, an ISS comprises the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine-3' (such as 5'-AACGTT-3').

In some embodiments, an ISS may comprise the sequence 5'-purine, T, C, G, pyrimidine, pyrimidine-3'.

In some embodiments, an ISS-containing polynucleotide is less than about any of the following lengths (in bases or base pairs): 10,000; 5,000; 2500; 2000; 1500; 1250; 1000; 750; 500; 300; 250; 200; 175; 150; 125; 100; 75; 50; 25; 10. In some embodiments, an ISS-containing polynucleotide is greater than about any of the following lengths (in bases or base pairs): 8; 10; 15; 20; 25; 30; 40; 50; 60; 75; 100; 125; 150; 175; 200; 250; 300; 350; 400; 500; 750; 1000; 2000; 5000; 7500; 10000; 20000; 50000. Alternately, the ISS can be any of a range of sizes having an upper limit of 10,000; 5,000; 2500; 2000; 1500; 1250; 1000; 750; 500; 300; 250; 200; 175; 150; 125; 100; 75; 50; 25; or 10 and an independently selected lower limit of 8; 10; 15; 20; 25; 30; 40; 50; 60; 75; 100; 125; 150; 175; 200; 250; 300; 350; 400; 500; 750; 1000; 2000; 5000; 7500, wherein the lower limit is less than the upper limit.

In some embodiments, the ISS comprises any of the following sequences:
GACGCTCC; GACGTCCC; GACGTTCC; GACGCCCC; AGCGTTCC;
AGCGCTCC; AGCGTCCC; AGCGCCCC; AACGTCCC; AACGCCCC;
AACGTTCC; AACGCTCC; GGCGTTCC; GGCGCTCC; GGCGTCCC;
GGCGCCCC; GACGCTCG; GACGTCCG; GACGCCCCG; GACGTTCCG;
AGCGCTCG; AGCGTTCCG; AGCGTCCG; AGCGCCCCG; AACGTCCG;
AACGCCCCG; AACGTTCCG; AACGCTCG; GGCGTTCCG; GGCGCTCG;
GGCGTCCG; GGCGCCCCG. In some embodiments, the immunomodulatory polynucleotide comprises the sequence 5'-TGACTGTGAACGTTTCGAGATGA-3' (SEQ ID NO:1).

In some embodiments, the ISS comprises any of the following sequences:
 GACGCU; GACGUC; GACGUU; GACGUT; GACGTU; AGCGUU; AGCGCU;
 AGCGUC; AGCGUT; AGCGTU; AACGUC; AACGUU; AACGCU; AACGUT;
 AACGTU; GGCGUU; GGCGCU; GGCGUC; GGCGUT; GGCGTU.

5 In some embodiments, the ISS comprises any of the following sequences:

GABGCTCC; GABGTCCC; GABGTTCC; GABGCCCC; AGBGTTCC;
 AGBGCTCC; AGBGTCCC; AGBGCCCC; AABGTCCC; AABGCCCC;
 AABGTTCC; AABGCTCC; GGBGTTCC; GGBGCTCC; GGBGTCCC;
 GGBGCCCC; GABGCTCG; GABGTCCG; GABGCCCCG; GABGTTCG;
 10 AGBGCTCG; AGBGTTCG; AGBGTCCG; AGBGCCCCG; AABGTCCG;
 AABGCCCCG; AABGTTCG; AABGCTCG; GGBGTTCG; GGBGCTCG;
 GGBGTCCG; GGBGCCCCG; GABGCTBG; GABGTCBG; GABGCCBG;
 GABGTTBG; AGBGCTBG; AGBGTTBG; AGBGTCBG; AGBGCCBG;
 AABGTCBG; AABGCCBG; AABGTTBG; AABGCTBG; GGBGTTBG;
 15 GGBGCTBG; GGBGTCBG; GGBGCCBG, where B is 5-bromocytosine.

In some embodiments, the ISS comprises any of the following sequences:

GABGCUCC; GABGUCCC; GABGUTCC; GABGTUCC; GABGUUCC;
 AGBGUUCC; AGBGTUCC; AGBGUTCC; AGBGCUCC; AGBGUCCC;
 AABGUCCC; AABGUUCC; AABGUTCC; AABGTUCC; AABGCUCC;
 20 GGBGUUCC; GGBGUTCC; GGBGTUCC; GGBGCUCC; GGBGUCCC;
 GABGCUCC; GABGUCCC; GABGUUCG; GABGUTC; GABGTUCG;
 AGBGCUCC; AGBGUUCG; AGBGUTC; AGBGTUCG; AGBGUCCC;
 AABGUCCC; AABGUUCG; AABGUTC; AABGTUCG; AABGCUCC;
 GGBGUUCG; GGBGUTC; GGBGTUCG; GGBGCUCC; GGBGUCCC;
 25 GABGCUBG; GABGUCBG; GABGUUBG; GABGUTBG; GABGTUBG;
 AGBGCUBG; AGBGUUBG; AGBGUCBG; AGBGUTBG; AGBGTUBG;
 AABGUCBG; AABGUUBG; AABGUTBG; AABGTUBG; AABGCUBG;

GGBGUUBG; GGBGUTBG; GGBGTUBG; GGBGCUBG; GGBGUCBG, where B is 5-bromocytosine.

In other embodiments, the ISS comprises any of the sequences:

5'-TGACCGTGAACGTTTCGAGATGA-3' (SEQ ID NO:2);

5 5'-TCATCTCGAACGTTCCACAGTCA-3' (SEQ ID NO:3);

5'-TGACTGTGAACGTTCCAGATGA-3' (SEQ ID NO:4);

5'-TCCATAACGTTTCGCCTAACGTTTCGTC-3' (SEQ ID NO:5);

5'-TGACTGTGAABGTTCCAGATGA-3' (SEQ ID NO:6), where B is 5-bromocytosine;

10 5'-TGACTGTGAABGTTTCGAGATGA-3' (SEQ ID NO:7), where B is 5-bromocytosine and

5'-TGACTGTGAABGTTBGAGATGA-3' (SEQ ID NO:8), where B is 5-bromocytosine.

An ISS and/or ISS-containing polynucleotide may contain modifications.

15 Modifications of ISS include any known in the art, but are not limited to, modifications of the 3'-OH or 5'-OH group, modifications of the nucleotide base, modifications of the sugar component, and modifications of the phosphate group. Various such modifications are described below.

20 An ISS may be single stranded or double stranded DNA, as well as single or double-stranded RNA or other modified polynucleotides. An ISS may or may not include one or more palindromic regions, which may be present in the motifs described above or may extend beyond the motif. An ISS may comprise additional flanking sequences, some of which are described herein. An ISS may contain naturally-occurring or modified, non-naturally occurring bases, and may contain
25 modified sugar, phosphate, and/or termini. For example, phosphate modifications include, but are not limited to, methyl phosphonate, phosphorothioate, phosphoramidate (bridging or non-bridging), phosphotriester and phosphorodithioate and may be used in any combination. Other non-phosphate linkages may also be

used. Preferably, polynucleotides of the present invention comprise phosphorothioate backbones. Sugar modifications known in the field, such as 2'-alkoxy-RNA analogs, 2'-amino-RNA analogs and 2'-alkoxy- or amino-RNA/DNA chimeras and others described herein, may also be made and combined with any phosphate modification.

5 Examples of base modifications include, but are not limited to, addition of an electron-withdrawing moiety to C-5 and/or C-6 of a cytosine of the ISS (e.g., 5-bromocytosine, 5-chlorocytosine, 5-fluorocytosine, 5-iodocytosine).

The ISS can be synthesized using techniques and nucleic acid synthesis equipment which are well known in the art including, but not limited to, enzymatic methods, chemical methods, and the degradation of larger polynucleotide sequences. See, for example, Ausubel et al. (1987); and Sambrook et al. (1989). When assembled enzymatically, the individual units can be ligated, for example, with a ligase such as T4 DNA or RNA ligase. U.S. Patent No. 5,124,246. Polynucleotide degradation can be accomplished through the exposure of an polynucleotide to a
10
15 nuclease, as exemplified in U.S. Patent No. 4,650,675.

The ISS can also be isolated using conventional polynucleotide isolation procedures. Such procedures include, but are not limited to, hybridization of probes to genomic or cDNA libraries and synthesis of particular native sequences by the polymerase chain reaction.

20 Circular ISS can be isolated, synthesized through recombinant methods, or chemically synthesized. Where the circular ISS is obtained through isolation or through recombinant methods, the ISS will preferably be a plasmid. The chemical synthesis of smaller circular oligonucleotides can be performed using any method described in the literature. See, for instance, Gao et al. (1995) *Nucleic Acids Res.* 23:2025-2029; and Wang et al. (1994) *Nucleic Acids Res.* 22:2326-2333.
25

The techniques for making polynucleotides and modified polynucleotides are known in the art. Naturally occurring DNA or RNA, containing phosphodiester linkages, is generally synthesized by sequentially coupling the appropriate nucleoside

phosphoramidite to the 5'-hydroxy group of the growing polynucleotide attached to a solid support at the 3'-end, followed by oxidation of the intermediate phosphite triester to a phosphate triester. Once the desired polynucleotide sequence has been synthesized, the polynucleotide is removed from the support, the phosphate triester groups are deprotected to phosphate diesters and the nucleoside bases are deprotected using aqueous ammonia or other bases. See, for example, Beaucage (1993) "Oligodeoxyribonucleotide Synthesis" in *Protocols for Oligonucleotides and Analogs, Synthesis and Properties* (Agrawal, ed.) Humana Press, Totowa, NJ; Warner et al. (1984) *DNA* 3:401 and U.S. Patent No. 4,458,066.

The ISS can also contain phosphate-modified polynucleotides. Synthesis of polynucleotides containing modified phosphate linkages or non-phosphate linkages is also known in the art. For a review, see Matteucci (1997) "Oligonucleotide Analogs: an Overview" in *Oligonucleotides as Therapeutic Agents*, (D.J. Chadwick and G. Cardew, ed.) John Wiley and Sons, New York, NY. The phosphorous derivative (or modified phosphate group) which can be attached to the sugar or sugar analog moiety in the polynucleotides of the present invention can be a monophosphate, diphosphate, triphosphate, alkylphosphonate, phosphorothioate, phosphorodithioate or the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and polynucleotides, *per se*, is also known and need not be described here in detail. Peyrottes et al. (1996) *Nucleic Acids Res.* 24:1841-1848; Chaturvedi et al. (1996) *Nucleic Acids Res.* 24:2318-2323; and Schultz et al. (1996) *Nucleic Acids Res.* 24:2966-2973. For example, synthesis of phosphorothioate polynucleotides is similar to that described above for naturally occurring polynucleotides except that the oxidation step is replaced by a sulfurization step (Zon (1993) "Oligonucleoside Phosphorothioates" in *Protocols for Oligonucleotides and Analogs, Synthesis and Properties* (Agrawal, ed.) Humana Press, pp. 165-190). Similarly the synthesis of other phosphate analogs, such as phosphotriester (Miller et al. (1971) *JACS* 93:6657-6665), non-bridging

phosphoramidates (Jager et al. (1988) *Biochem.* 27:7247-7246), N3' to P5' phosphoramidates (Nelson et al. (1997) *JOC* 62:7278-7287) and phosphorodithioates (U.S. Patent No. 5,453,496) has also been described. Other non-phosphorous based modified polynucleotides can also be used (Stirchak et al. (1989) *Nucleic Acids Res.* 17:6129-6141). Polynucleotides with phosphorothioate backbones can be more immunogenic than those with phosphodiester backbones and appear to be more resistant to degradation after injection into the host. Braun et al. (1988) *J. Immunol.* 141:2084-2089; and Latimer et al. (1995) *Mol. Immunol.* 32:1057-1064.

ISS-containing polynucleotides used in the invention can comprise ribonucleotides (containing ribose as the only or principal sugar component), deoxyribonucleotides (containing deoxyribose as the principal sugar component), or, as is known in the art, modified sugars or sugar analogs can be incorporated in the ISS. Thus, in addition to ribose and deoxyribose, the sugar moiety can be pentose, deoxypentose, hexose, deoxyhexose, glucose, arabinose, xylose, lyxose, and a sugar "analog" cyclopentyl group. The sugar can be in pyranosyl or in a furanosyl form. In the ISS, the sugar moiety is preferably the furanoside of ribose, deoxyribose, arabinose or 2'-O-alkylribose, and the sugar can be attached to the respective heterocyclic bases either in α or β anomeric configuration. Sugar modifications include, but are not limited to, 2'-alkoxy-RNA analogs, 2'-amino-RNA analogs and 2'-alkoxy- or amino-RNA/DNA chimeras. The preparation of these sugars or sugar analogs and the respective "nucleosides" wherein such sugars or analogs are attached to a heterocyclic base (nucleic acid base) *per se* is known, and need not be described here, except to the extent such preparation can pertain to any specific example. Sugar modifications may also be made and combined with any phosphate modification in the preparation of an ISS.

The heterocyclic bases, or nucleic acid bases, which are incorporated in the ISS can be the naturally-occurring principal purine and pyrimidine bases, (namely

uracil or thymine, cytosine, adenine and guanine, as mentioned above), as well as naturally-occurring and synthetic modifications of said principal bases.

Those skilled in the art will recognize that a large number of "synthetic" non-natural nucleosides comprising various heterocyclic bases and various sugar
5 moieties (and sugar analogs) are available in the art, and that as long as other criteria of the present invention are satisfied, the ISS can include one or several heterocyclic bases other than the principal five base components of naturally-occurring nucleic acids. Preferably, however, the heterocyclic base in the ISS includes, but is not limited to, uracil-5-yl, cytosin-5-yl, adenin-7-yl, adenin-8-yl, guanin-7-yl,
10 guanin-8-yl, 4-aminopyrrolo (2.3-d) pyrimidin-5-yl, 2-amino-4-oxopyrrolo (2,3-d) pyrimidin-5-yl, 2-amino-4-oxopyrrolo (2.3-d) pyrimidin-3-yl groups, where the purines are attached to the sugar moiety of the ISS via the 9-position, the pyrimidines via the 1-position, the pyrrolopyrimidines via the 7-position and the pyrazolopyrimidines via the 1-position.

The ISS may comprise at least one modified base as described, for example, in the commonly owned international application WO 99/62923. As used herein, the term "modified base" is synonymous with "base analog", for example, "modified cytosine" is synonymous with "cytosine analog." Similarly, "modified" nucleosides or nucleotides are herein defined as being synonymous with nucleoside or nucleotide
15 "analogs." Examples of base modifications include, but are not limited to, addition of an electron-withdrawing moiety to C-5 and/or C-6 of a cytosine of the ISS. Preferably, the electron-withdrawing moiety is a halogen. Such modified cytosines can include, but are not limited to, azacytosine, 5-bromocytosine, bromouracil, 5-chlorocytosine, chlorinated cytosine, cyclocytosine, cytosine arabinoside, 5-
20 fluorocytosine, fluoropyrimidine, fluorouracil, 5,6-dihydrocytosine, 5-iodocytosine, hydroxyurea, iodouracil, 5-nitrocytosine, uracil, and any other pyrimidine analog or modified pyrimidine.

The preparation of base-modified nucleosides, and the synthesis of modified polynucleotides using said base-modified nucleosides as precursors, has been described, for example, in U.S. Patents 4,910,300, 4,948,882, and 5,093,232. These base-modified nucleosides have been designed so that they can be incorporated by chemical synthesis into either terminal or internal positions of an polynucleotide. Such base-modified nucleosides, present at either terminal or internal positions of an polynucleotide, can serve as sites for attachment of a peptide or other antigen. Nucleosides modified in their sugar moiety have also been described (including, but not limited to, *e.g.*, U.S. Patents 4,849,513, 5,015,733, 5,118,800, 5,118,802) and can be used similarly.

The ISS used in the methods of the invention may be produced as ISS-microcarrier complexes. ISS-microcarrier complexes comprise an ISS-containing polynucleotide bound to a microcarrier (MC). ISS-MC complexes comprise an ISS bound to the surface of a microcarrier (*i.e.*, the ISS is not encapsulated in the MC), adsorbed within a microcarrier (*e.g.*, adsorbed to PLGA beads), or encapsulated within a MC (*e.g.*, incorporated within liposomes).

ISS-containing oligonucleotides bound to microparticles (SEPHAROSE® beads) have previously been shown to have immunostimulatory activity *in vitro* (Liang et al., (1996), *J. Clin. Invest.* 98:1119-1129). However, recent results show that ISS-containing oligonucleotides bound to gold, latex and magnetic particles are not active in stimulating proliferation of 7TD1 cells, which proliferate in response to ISS-containing oligonucleotides (Manzel et al., (1999), *Antisense Nucl. Acid Drug Dev.* 9:459-464).

Microcarriers are not soluble in pure water, and are less than about 50-60 μm in size, preferably less than about 10 μm in size, more preferably from about 10 nm to about 10 μm , 25 nm to about 5 μm , 50 nm to about 4.5 μm or 1.0 μm to about 2.0 μm in size. Microcarriers may be any shape, such as spherical, ellipsoidal, rod-shaped, and the like, although spherical microcarriers are normally preferred.

Preferred microcarriers have sizes of or about 50 nm, 200 nm, 1 μm , 1.2 μm , 1.4 μm , 1.5 μm , 1.6 μm , 1.8 μm , 2.0 μm , 2.5 μm or 4.5 μm . The “size” of a microcarrier is generally the “design size” or intended size of the particles stated by the manufacturer. Size may be a directly measured dimension, such as average or maximum diameter, or may be determined by an indirect assay such as a filtration screening assay. Direct measurement of microcarrier size is typically carried out by microscopy, generally light microscopy or scanning electron microscopy (SEM), in comparison with particles of known size or by reference to a micrometer. As minor variations in size arise during the manufacturing process, microcarriers are considered to be of a stated size if measurements show the microcarriers are \pm about 5-10% of the stated measurement. Size characteristics may also be determined by dynamic light scattering. Alternately, microcarrier size may be determined by filtration screening assays. A microcarrier is less than a stated size if at least 97% of the particles pass through a “screen-type” filter (*i.e.*, a filter in which retained particles are on the surface of the filter, such as polycarbonate or polyethersulfone filters, as opposed to a “depth filter” in which retained particles lodge within the filter) of the stated size. A microcarrier is larger than a stated size if at least about 97% of the microcarrier particles are retained by a screen-type filter of the stated size. Thus, at least about 97% microcarriers of about 10 μm to about 10 nm in size pass through a 10 μm pore screen filter and are retained by a 10 nm screen filter.

As above discussion indicates, reference to a size or size range for a microcarrier implicitly includes approximate variations and approximations of the stated size and/or size range. This is reflected by use of the term “about” when referring to a size and/or size range, and reference to a size or size range without reference to “about” does not mean that the size and/or size range is exact.

Microcarriers may be solid phase (*e.g.*, polystyrene beads) or liquid phase (*e.g.*, liposomes, micelles, or oil droplets in an oil and water emulsion). Liquid phase microcarriers include liposomes, micelles, oil droplets and other lipid or oil-based

particles. One preferred liquid phase microcarrier is oil droplets within an oil-in-water emulsion. Preferably, oil-in-water emulsions used as microcarriers comprise biocompatible substituents such as squalene. Liquid phase microcarriers are normally considered nonbiodegradable, but may be biodegradable liquid phase microcarriers may be produced by incorporation of one or more biodegradable polymers in the liquid microcarrier formulation. In one preferred embodiment, the microcarrier is oil droplets in an oil-in-water emulsion prepared by emulsification of squalene, sorbitan trioleate, TWEEN 80® in an aqueous pH buffer.

Solid phase microcarriers for use in ISS-microcarrier complexes may be made from biodegradable materials or nonbiodegradable materials, and may include or exclude agarose or modified agarose microcarriers. Useful solid phase biodegradable microcarriers include, but are not limited to: biodegradable polyesters, such as poly(lactic acid), poly(glycolic acid), and copolymers (including block copolymers) thereof, as well as block copolymers of poly(lactic acid) and poly(ethylene glycol); polyorthoesters such as polymers based on 3,9-diethylidene-2,4,8,10-tetraoxaspiro[5.5]undecane (DETOSU); polyanhydrides such as poly(anhydride) polymers based on sebacic acid, *p*-(carboxyphenoxy)propane, or *p*-(carboxyphenoxy)hexane; polyanhydride imides, such as polyanhydride polymers based on sebacic acid-derived monomers incorporating amino acids (*i.e.*, linked to sebacic acid by imide bonds through the amino-terminal nitrogen) such as glycine or alanine; polyanhydride esters; polyphosphazenes, especially poly(phosphazenes) which contain hydrolysis-sensitive ester groups which can catalyze degradation of the polymer backbone through generation of carboxylic acid groups (Schacht et al. (1996) *Biotechnol. Bioeng.* 1996:102); and polyamides such as poly(lactic acid-co-lysine). A wide variety of nonbiodegradable materials suitable for manufacturing microcarriers are also known, including, but not limited to polystyrene, polyethylene, latex, gold, and ferromagnetic or paramagnetic materials. Solid phase microcarriers may be covalently modified to incorporate one or more moieties for use in linking the

ISS, for example by addition of amine groups for covalent linking using amine-reactive crosslinkers.

The ISS-microcarrier complexes of the invention may be covalently or non-covalently linked. Covalently linked ISS-MC complexes may be directly linked or be
5 linked by a crosslinking moiety of one or more atoms (typically the residue of a crosslinking agent). The ISS may be modified to allow or augment binding to the MC (e.g., by incorporation of a free sulfhydryl for covalent crosslinking or addition of a hydrophobic moieties such as lipids, steroids, sterols such as cholesterol, and
10 terpenes, for hydrophobic bonding), although unmodified ISS may be used for formation of non-covalent ISS-MC complex formation by electrostatic interaction or by base pairing (e.g., by base pairing at least one portion of the ISS with a complementary oligonucleotide bound to the microcarrier). ISS-containing polynucleotides may be linked to solid phase microcarriers or other chemical
15 moieties to facilitate ISS-MC complex formation using conventional technology known in the art, such as use of available heterobifunctional crosslinkers (e.g., succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate or its sulfo-derivatives for covalently linking an amine-derivatized microcarrier and an ISS
20 modified to contain a free sulfhydryl) or by addition of compounds such as cholesterol (e.g., by the method of Godard et al. (1995) *Eur. J. Biochem.* 232:404-410) to facilitate binding to hydrophobic microcarriers such as oil droplets in oil-in-water emulsions. Alternatively, modified nucleosides or nucleotides, such as are
25 known in the art, can be incorporated at either terminus, or at internal positions in the ISS. These can contain blocked functional groups which, when deblocked, are reactive with a variety of functional groups which can be present on, or attached to, the microcarrier or a moiety which would facilitate binding to a microcarrier. Certain
embodiments of noncovalently linked ISS-MC complexes utilize a binding pair (e.g., an antibody and its cognate antigen or biotin and streptavidin or avidin), where one member of the binding pair is bound to the ISS and the microcarrier is derivatized

with the other member of the binding pair (*e.g.*, a biotinylated ISS and a streptavidin-derivatized microcarrier may be combined to form a noncovalently linked ISS-MC complex).

Non-covalent ISS-MC complexes bound by electrostatic binding typically exploit the highly negative charge of the polynucleotide backbone. Accordingly, microcarriers for use in non-covalently bound ISS-MC complexes are generally positively charged at physiological pH (*e.g.*, about pH 6.8-7.4). The microcarrier may intrinsically possess a positive charge, but microcarriers made from compounds not normally possessing a positive charge may be derivatized or otherwise modified to become positively charged. For example, the polymer used to make the microcarrier may be derivatized to add positively charged groups, such as primary amines. Alternately, positively charged compounds may be incorporated in the formulation of the microcarrier during manufacture (*e.g.*, positively charged surfactants may be used during the manufacture of poly(lactic acid)/poly(glycolic acid) copolymers to confer a positive charge on the resulting microcarrier particles.

Solid phase microspheres are prepared using techniques known in the art. For example, they can be prepared by emulsion-solvent extraction/evaporation technique. Generally, in this technique, biodegradable polymers such as polyanhydrides, poly(alkyl- α -cyanoacrylates) and poly(α -hydroxy esters), for example, poly(lactic acid), poly(glycolic acid), poly(D,L-lactic-co-glycolic acid) and poly(caprolactone), are dissolved in a suitable organic solvent, such as methylene chloride, to constitute the dispersed phase (DP) of emulsion. DP is emulsified by high-speed homogenization into excess volume of aqueous continuous phase (CP) that contains a dissolved surfactant, for example, polyvinylalcohol (PVA) or polyvinylpyrrolidone (PVP). Surfactant in CP is to ensure the formation of discrete and suitably-sized emulsion droplet. The organic solvent is then extracted into the CP and subsequently evaporated by raising the system temperature. The solid microparticles are then

separated by centrifugation or filtration, and dried, for example, by lyophilization or application of vacuum, before storing at 4 °C.

Generally, to prepare cationic microspheres, cationic lipids or polymers, for example, 1,2-dioleoyl-1,2,3-trimethylammonio propane (DOTAP),
5 cetyltrimethylammonium bromide (CTAB) or polylysine, are added either to DP or CP, as per their solubility in these phases.

Physico-chemical characteristics such as mean size, size distribution and surface charge of dried microspheres may be determined. Size characteristics are determined, for example, by dynamic light scattering technique and the surface
10 charge was determined by measuring the zeta potential.

Generally, ISS-containing polynucleotides can be adsorbed onto the cationic microspheres by overnight aqueous incubation of ISS and the particles at 4 °C. Microspheres are characterized for size and surface charge before and after ISS
15 association. Selected batches may then be evaluated for activity as described herein.

Administration

An ISS-containing polynucleotide may be administered before, during and/or after exposure to a papillomavirus. An ISS polynucleotide may also be administered before, during and/or after infection by a papillomavirus. An ISS-containing
20 polynucleotide may also be administered before or after onset of symptoms of papillomavirus infection. An ISS-containing polynucleotide may also be administered before the development of papillomavirus-associated carcinoma (*i.e.*, a pre-cancerous state). Accordingly, administration of ISS-containing polynucleotide may be at various times with respect to exposure to, infection by and/or onset of
25 symptoms of infection by papillomavirus. Further, there may be one or more administrations. If the ISS-containing polynucleotide is administered on multiple occasions, the ISS may be administered on any schedule selected by the clinician, such as daily, every other day, every three days, every four days, every five days,

every six days, weekly, biweekly, monthly or at ever longer intervals (which may or may not remain the same during the course of treatment). Where multiple administrations are given, the ISS-containing polynucleotide may be given in 2, 3, 4, 5, 6, 7, 8, 9, 10 or more separate administrations.

5 When ISS-containing polynucleotide is administered to an individual at risk of exposure to virus (*i.e.*, before infection), ISS-containing polynucleotide is preferably administered less than about 14 days before exposure to virus, preferably less than about 10 days before exposure to virus, more preferably less than about 7 days before exposure to virus, even more preferably less than about 5 days before exposure to virus. In some embodiments, ISS-containing polynucleotide is administered about 3 days before exposure to virus.

10 In a further embodiment, the ISS-containing polynucleotide is administered after exposure to or infection by a papillomavirus, but prior to appearance of symptoms. This embodiment is particularly relevant with respect to HPV since years can elapse between exposure to papillomavirus and possible progression to carcinoma. Preferably, the ISS-containing polynucleotide is administered less than about three days after exposure, more preferably less than about one day, 12 hours, six hours or two hours after exposure, if the time of exposure is known or suspected.

15 In another embodiment, the ISS-containing polynucleotide is administered after appearance of at least one symptom of papillomavirus infection. Preferably, ISS-containing polynucleotide is administered within about 28, 21, 14, 7, 5 or 3 days following appearance of a symptom of papillomavirus infection. However, some infected individuals exhibiting symptoms will already have undertaken one or more courses of treatment with another therapy. In such individuals, or in individuals who failed to appreciate the import of their symptoms, the ISS-containing polynucleotide may be administered at any point following infection.

20 In another embodiment, the ISS-containing polynucleotide is administered after the appearance of at least one symptom of papillomavirus infection and

preferably before the development of carcinoma. The staging of abnormal cellular growth (dysplasia) may be accomplished by any of the methods known in art, including, but not limited to, Pap smears, local biopsies, and *in situ* hybridization.

5 Additionally, treatments employing an ISS-containing polynucleotide may also be employed in conjunction with other treatments or as 'second line' treatments employed after failure of a 'first line' treatment (*e.g.*, ISS-containing polynucleotide therapy may be employed in conjunction with physical removal and/or cryogenic treatment of papillomavirus-induced lesions).

10 ISS polynucleotides may be formulated in any form known in the art, such as dry powder, semi-solid or liquid formulations. For parenteral administration ISS polynucleotides preferably administered in a liquid formulation, although solid or semi-solid formulations may also be acceptable, particularly where the ISS polynucleotide is formulated in a slow release depot form. ISS polynucleotides are generally formulated in liquid or dry powder form for topical administration,
15 although semi-solid formulations may occasionally be useful.

ISS polynucleotide formulations may contain additional components such as salts, buffers, bulking agents, osmolytes, antioxidants, detergents, surfactants and other pharmaceutically-acceptable excipients as are known in the art. Generally, liquid ISS polynucleotide formulations made in USP water for injection and are
20 sterile, isotonic and pH buffered to a physiologically-acceptable pH, such as about pH 6.8 to 7.5.

ISS-containing polynucleotides may be formulated in delivery vehicles such as liposomes, oil/water emulsion or slow release depot formulations. Methods of formulating polynucleotides in such forms are well known in the art.

25 ISS-containing polynucleotide formulations may also include or exclude immunomodulatory agents such as adjuvants and immunostimulatory cytokines, which are well known in the art.

A suitable dosage range or effective amount is one that provides the desired reduction of symptoms and/or suppression of viral infection and depends on a number of factors, including the particular papillomavirus, ISS sequence of the polynucleotide, molecular weight of the polynucleotide and route of administration.

5 Dosages are generally selected by the physician or other health care professional in accordance with a variety of parameters known in the art, such as severity of symptoms, history of the patient and the like. Generally, for an ISS-containing polynucleotide of about 20 bases, a dosage range may be selected from, for example, an independently selected lower limit such as about 0.1, 0.25, 0.5, 1, 2, 5, 10, 20, 30
10 40, 50 60, 80, 100, 200, 300, 400 or 500 µg/kg up to an independently selected upper limit, greater than the lower limit, of about 60, 80, 100, 200, 300, 400, 500, 750, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000 or 10,000 µg/kg. For example, a dose may be about any of the following: 0.1 to 100 µg/kg, 0.1 to 50 µg/kg, 0.1 to 25 µg/kg, 0.1 to 10 µg/kg, 1 to 500 µg/kg, 100 to 400 µg/kg, 200 to 300
15 µg/kg, 1 to 100 µg/kg, 100 to 200 µg/kg, 300 to 400 µg/kg, 400 to 500 µg/kg, 500 to 1000 µg/kg, 500 to 5000 µg/kg, or 500 to 10,000 µg/kg. Generally, parenteral routes of administration require higher doses of ISS compared to more direct application to infected tissue, as do ISS-containing polynucleotides of increasing length.

Polynucleotides comprising an ISS may be administered by systemic (*e.g.*,
20 parenteral) or local (*e.g.*, topical or intralesional injection) administration.

In one embodiment, the ISS-containing polynucleotide(s) is topically administered. Topical administration may be at the site of infection (*e.g.*, genital region in the case of mucosal papillomavirus), it may be at a site of a symptom (*e.g.*, a papilloma lesion) or it may be at the site of possible exposure to papillomavirus
25 (*e.g.*, genital region).

In another embodiment, the ISS-containing polynucleotide(s) is injected locally into the area of lesion(s). Intralesional injection may be at the site of infection (*e.g.*, genital region in the case of mucosal papillomavirus), site of dysplasia (*e.g.*,

epithelium in the genital region) or it may be at a site of a symptom (*e.g.*, a papilloma lesion).

In other embodiments, the ISS-containing polynucleotide is administered parenterally. Parenteral routes of administration include, but are not limited to, transdermal, transmucosal, nasopharyngeal, pulmonary and direct injection. Parenteral administration by injection may be by any parenteral injection route, including, but not limited to, intravenous (IV), intraperitoneal (IP), intramuscular (IM), subcutaneous (SC) and intradermal (ID) routes. Transdermal and transmucosal administration may be accomplished by, for example, inclusion of a carrier (*e.g.*, dimethylsulfoxide, DMSO), by application of electrical impulses (*e.g.*, iontophoresis) or a combination thereof. A variety of devices are available for transdermal administration which may be used in accordance with the invention.

Nasopharyngeal and pulmonary routes of administration include, but are not limited to, intranasal, inhalation, transbronchial and transalveolar routes. The ISS-containing polynucleotide may thus be administered by inhalation of aerosols, atomized liquids or powders. Devices suitable for administration by inhalation of ISS-containing compositions include, but are not limited to, nebulizers, atomizers, vaporizers, and metered-dose inhalers. Nebulizers, atomizers, vaporizers and metered-dose inhalers filled with or employing reservoirs containing formulations comprising the ISS-containing polynucleotide(s) are among a variety of devices suitable for use in inhalation delivery of the ISS-containing polynucleotide(s). Other methods of delivering to respiratory mucosa include delivery of liquid formulations, such as by nose drops.

IV, IP, IM and ID administration may be by bolus or infusion administration. For SC administration, administration may be by bolus, infusion or by implantable device, such as an implantable minipump (*e.g.*, osmotic or mechanical minipump) or slow release implant. The ISS polynucleotide(s) may also be delivered in a slow release formulation adapted for IV, IP, IM, ID or SC administration. Administration

by inhalation is preferably accomplished in discrete doses (e.g., via a metered dose inhaler), although delivery similar to an infusion may be accomplished through use of a nebulizer. Administration via the transdermal and transmucosal routes may be continuous or pulsatile.

5

Assessment

In some embodiments, administration of an ISS-containing polynucleotide results in prevention, palliation and/or improvement in one or more symptoms of papillomavirus infection. The exact form of prevention, palliation or improvement will depend on the particular *papillomavirinae* type and the symptoms experienced by the individual but includes reduction in size and/or duration of lesions and/or warts, reduction in symptoms of papillomavirus infection or reduction in frequency or number of recurrent lesions. In some embodiments, administration of an ISS-containing polynucleotide results in a reduction in viral titer (a reduction of which indicates suppression of viral infection). In other embodiments, the number of warts is reduced. In other embodiments, viral infection is suppressed, which may be indicated by any one or more of a number of parameters, including, but not limited to, extent of one or more symptoms and viral titer. In other embodiments, recurrence, which is generally indicated by appearance of one or more symptoms associated with infection, is reduced.

10

15

20

Symptoms of infection may be assessed before or after administration of ISS-containing polynucleotide by the individual or the clinician. As will be apparent to one of skill in the art, the symptoms will vary depending on the particular papillomavirus (e.g., cutaneous or mucosal, high-risk or low-risk) and the site of the symptoms (e.g., genital region, oral cavity, respiratory tract, skin, etc.). Symptoms of papillomavirus infection can include papilloma lesions on cutaneous and/or mucosal membranes, thickening of epithelial layer, nuclear changes such as enlargement, hyperchromasia, and/or pyknosis. Characteristics of papillomavirus lesions can

25

include localized epithelial hyperplasia with a defined boundary, intact basement membrane, and differentiated epithelium. Additional characteristics may include koilocytosis, large perinuclear cavitation with irregular edges and dense cytoplasm in the area surrounding the cavity.

5 Viral titer may be assessed in biological samples using standard methods of the art. Levels of viral nucleic acid may be assessed by isolating nucleic acid from the sample and/or performing PCR analysis using virus specific primers or blot analysis using a viral polynucleotide sequence as a probe. The PCR analysis may be quantitative using PCR technology known in the art. Another method is to perform
10 *in situ* hybridization with virus-specific probes. Another assay measures infectious units, such as infectious center assay (ICA). Extent or amount of viral particles may be measured from any infected area, such as infected tissue or mucosal discharge. When the sample is a liquid, viral titer is calculated in some indication of number or amount of virus or virus particles (*e.g.*, infectious particles, plaque forming units, infectious doses, or median tissue culture infectious doses (TCID 50)) per unit
15 volume. In solid samples, such as a tissue sample, viral titer is calculated in virus particles per unit weight. Reduction is indicated by comparing viral titer to viral titer measured at an earlier time point, and/or comparing to an estimated titer (based, for example, on animal or clinical studies) that represents untreated infection.

20 Abnormal cell growth in the cervix may be examined and classified into various stages according to procedures commonly practiced by clinicians. Samples obtained by methods such as Pap smears or local biopsy may be examined for morphological abnormalities. Abnormal cell growth in the cervix may be classified as either low-grade squamous intraepithelial lesion (SIL) or high-grade SIL. The
25 category of low-grade SIL includes HPV infection and cervical intraepithelial neoplasia (CIN) 1 while high-grade SIL includes CIN 2 and 3 when the entire thickness of the epithelium is replaced by abnormal cells. This classification scheme is equivalent to a previous classification scheme in which the different stages ranged

from cervical intraepithelial neoplasia (CIN) 1 (mild dysplasia) to CIN 2 (moderate dysplasia) to CIN 3 (carcinoma *in situ*). Individuals with HPV infection but have not progressed to CIN1 are suitable candidates for the administration of ISS to prevent and/or reduce the chances of progression to carcinoma.

5

Kits of the Invention

The invention provides kits for carrying out the methods of the invention. Accordingly, a variety of kits are provided. The kits may be used for any one or more of the following (and, accordingly, may contain instructions for any one or more of the following uses): preventing symptoms of *papillomavirinae* infection in an individual who has been exposed to *papillomavirinae*; preventing symptoms of *papillomavirinae* infection in an individual at risk of being exposed to *papillomavirinae*; reducing severity of a symptom of *papillomavirinae* infection in an individual infected with *papillomavirinae*; reducing recurrence of a symptom of *papillomavirinae* infection in an individual infected with *papillomavirinae*; delaying development of a *papillomavirinae* infection and/or a symptom of *papillomavirinae* infection in an individual infected or at risk of being infected with *papillomavirinae*; reducing duration of a *papillomavirinae* infection in an individual infected or at risk of being infected with *papillomavirinae*. As is understood in the art, any one or more of these uses would be included in instructions directed to treating or preventing *papillomavirinae* infection.

The kits of the invention comprise one or more containers comprising an ISS-containing polynucleotide and a set of instructions, generally written instructions although electronic storage media (*e.g.*, magnetic diskette or optical disk) containing instructions are also acceptable, relating to the use and dosage of the ISS-containing polynucleotide for the intended treatment (*e.g.*, preventing one or more symptoms of *papillomavirinae* infection in an individual who has been exposed to *papillomavirinae*; preventing one or more symptoms of *papillomavirinae* infection in

an individual who is at risk of being exposed to *papillomavirinae*; reducing severity of a symptom of *papillomavirinae* infection in an individual infected with *papillomavirinae*; and/or reducing recurrence of one or more symptoms of *papillomavirinae* infection in an individual infected with *papillomavirinae*). The instructions included with the kit generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. The containers of ISS may be unit doses, bulk packages (e.g., multi-dose packages) or sub-unit doses.

The kits of the invention do not include any packages or containers which include viral antigens from the *papillomavirinae* the kit is intended to be used to treat. Accordingly, neither the container comprising the ISS-containing polynucleotide nor any other containers in the kit contain *papillomavirinae* viral antigens.

The ISS component of the kit may be packaged in any convenient, appropriate packaging. For example, if the ISS is a freeze-dried formulation, a vial with a resilient stopper is normally used, so that the drug may be easily reconstituted by injecting fluid through the resilient stopper. Ampoules with non-resilient, removable closures (e.g., sealed glass) or resilient stoppers are most conveniently used for injectable forms of ISS. Also, prefilled syringes may be used when the kit is supplied with a liquid formulation of the ISS-containing polynucleotide. The kit may contain the ISS in an ointment for topical formulation in appropriate packaging. Also contemplated are packages for use in combination with a specific device, such as an inhaler, nasal administration device (e.g., an atomizer), transdermal administration device, or an infusion device such as a minipump.

As stated above, any ISS-containing polynucleotide described herein may be used, such as, for example, any polynucleotide comprising any of the following ISS: the sequence 5'-C, G, pyrimidine, pyrimidine, C, G-3', the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine, C, G-3', the sequence 5'-purine, purine, C, G, pyrimidine, pyrimidine, C, C-3'; the sequence SEQ ID NO: 1; the sequence 5'-purine, purine, B, G, pyrimidine, pyrimidine-3' wherein B is 5-bromocytosine or the

sequence 5'-purine, purine, B, G, pyrimidine, pyrimidine, C, G-3' wherein B is 5-bromocytosine.

The following Examples are provided to illustrate, but not limit, the invention.

5

EXAMPLES

Example 1: Treatment of Canine Oral Papilloma with ISS

A model of canine oral papilloma was used to test the efficacy of ISS on papilloma. Beagle puppies were inoculated in the bucal mucosa with canine papillomavirus and developing papilloma lesions were monitored daily. Four groups of seven dogs each were treated with differing amount of ISS oligonucleotide (5'-TGACTGTGAACGTTTCGAGATGA-3' (SEQ ID NO:1), phosphorothioate backbone). One group received 50 µg ISS twice a week, another group received 500 µg ISS twice a week, the third group received 500 µg ISS one time only at the first signs of papilloma lesion development (injected within the papilloma lesion) and the fourth group (control group) received PBS twice a week. All dogs were monitored daily for the development of lesions and the time to regression.

The results are shown in Figure 1. Dogs that received a one time treatment of 500 µg ISS at the first signs of papilloma lesion showed a higher average rate of lesion regression than untreated dogs, although the ranges for both groups overlapped. Untreated dogs took an average of 29.1 days for rapid regression while dogs treated with 500 µg ISS at the first signs of papilloma took an average of 25.1 days for rapid regression.

The other treatment groups did not show a marked difference in regression time. This model offers a short window of time in which regression of warts can be observed. In dogs, warts caused by canine papillomaviruses can spontaneously regress. Injection of ISS in the papillomas when papillomas first appear appears to

enhance the time of lesion regression as compared to the time of spontaneous lesion regression.

Example 2: Treatment of cutaneous papillomatosis in a rabbit model by ISS

5 Rabbits were initially the first animals in which papillomavirus infection was described in 1933 by Shope. Shope recognized the cottontail rabbit papillomavirus (CRPV) as the etiological agent for cutaneous papillomatosis in the cottontail rabbit (Howley, P., Chapter 65, *Fields Virology*, Vol. 2, Third Edition, Lippincott-Raven publishers).

10 In this model of papilloma, New Zealand White rabbits of both genders were quarantined for 14 days, those animals remaining healthy were cleared for use in the experiment. 15 rabbits were each inoculated with a high dose of CRPV at two different sites (one on each side fo the animal) and a low dose of CRPV at two different sites (one on each side fo the animal) for a total of four inoculation sites in
15 each rabbit. The animals were then separated into three groups of five animals each, groups A, B, and C.

Group A received 50 µg intradermal injections of ISS oligonucleotide (5'-TGACTGTGAACGTTTCGAGATGA-3' (SEQ ID NO:1), phosphorothioate backbone) into the site of CRPV inoculation (site of the papilloma lesion at later time
20 points) at Day 1 (one day following inoculation with CRPV) and Day 21 on the left side and at Day 14 and Day 35 on the right side. Groups B and C received intradermal injections of 500 µg of the ISS and phosphate-buffered saline (vehicle), respectively, into the site of CRPV inoculation (site of the papilloma lesion at later
time points) on the same schedule.

25 Papilloma development was quantitated by finding the geometric mean diameter (GMD) of each papilloma lesion. GMD was calculated from measurements of the length, width and height of the papilloma lesions. Measurements were made weekly.

Results are summarized in Figure 2. Panel A shows GMD for the left side, high CRPV dose lesions (treatment on Day 1 and 14). Panel B shows GMD for the left side, low CRPV dose lesions (treatment on Day 14 and 35). Panel C shows average GMD for the right side, high CRPV dose lesions (treatment on Day 1 and 14). Panel D shows average GMD for the right side, low CRPV dose lesions (treatment on Day 14 and 35).

In another experiment, a mutant of CRPV which induces small papillomas, CRPV-E8m, was used to induce papillomas on five rabbit treatment groups (five rabbits per group). In each animal, papillomas on the left side of the animal received treatments and papillomas on the right side were untreated. Four of the treatment groups received doses between 100 µg and 2000 µg of ISS as intradermal injections per papilloma at several treatment regimes and the fifth group received injections of PBS as control, as outlined below.

<u>Group</u>	<u>Left Side Treatment</u>
A	ISS; 100 µg/injection; 3 times/week from days 47 - 86
B	ISS; 100 µg/injection; 1 time/week from days 47 - 86
C	ISS; 500 µg/injection; 1 time/week from days 47 - 86
D	ISS; 2000 µg/injection; weeks 7 and 10
E	PBS; 100 µl/injection; 3 times/week from days 47 - 86

Four papillomas, initiated with CRPV-E8m plasmid DNA, were established on each rabbit. Skin at the site of papilloma initiation was made hyperplastic using a mixture of turpentine and acetone prior to viral DNA administration. The size of papillomas was measured (three dimensions, in mm) and the GMD calculated for each papilloma.

In this experiment, a number of viral DNA challenged sites failed to generate any papillomas. Minimal differences were found in the papilloma growth rates of the treated versus untreated papillomas for Treatment Groups A, B, D and E. Results

from Treatment Group C are depicted in Figure 3 and demonstrate a reduction in the size of the ISS treated papillomas compared to untreated papillomas.

5 The present invention has been detailed both by direct description and by example. Equivalents and modifications of the present invention will be apparent to those skilled in the art, and are encompassed within the scope of the invention.

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
224